

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Organic and inorganic carbon in the topsoil of the Mongolian and Tibetan grasslands: pattern, control and implications

Y. Shi¹, F. Baumann², Y. Ma¹, C. Song^{1,*}, P. Kühn², T. Scholten², and J.-S. He³

¹Department of Ecology, College of Urban and Environmental Sciences, and Key Laboratory for Earth Surface Processes of the Ministry of Education, Peking University, 5 Yiheyuan Rd., 100871 Beijing, China

²Department of Geoscience, Physical Geography and Soil Science, University of Tuebingen, Ruemelinstrasse 19–23, 72070 Tuebingen, Germany

³Key Laboratory of Adaptation and Evolution of Plateau Biota, Northwest Institute of Plateau Biology, Chinese Academy of Sciences, 23 Xinning Rd., 810008 Xining, China

*now at: Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, KS 66045, USA

Received: 11 January 2012 – Accepted: 4 February 2012 – Published: 15 February 2012

Correspondence to: J.-S. He (jshe@pku.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

1869

Abstract

Soil carbon (C) is the largest C pool in terrestrial biosphere and includes both inorganic and organic components. Studying patterns and controls of soil C help us to understand and estimate potential responses of soil C to global change in the future. Here we analyzed topsoil data of 81 sites obtained from a regional survey across grasslands in the Inner Mongolia and on the Tibetan Plateau during 2006–2007, attempting to find the patterns and controls of soil inorganic carbon (SIC) and soil organic carbon (SOC). The average of SIC and SOC in the topsoil (0–20 cm) across the study region were 0.38 % and 3.63 %, ranging between 0.00–2.92 % and 0.32–26.17 %, respectively. Both SIC and SOC in the topsoil of the Tibetan grasslands (0.51 % and 5.24 %, respectively) were higher than those of the Inner Mongolian grasslands (0.21 % and 1.61 %). Regression tree analyses showed that the spatial pattern of SIC and SOC were controlled by different factors. Chemical and physical processes of soil formation drive the spatial pattern of SIC, while biotic processes drive the spatial pattern of SOC. SIC was controlled by soil acidification and other processes depending on soil pH. Vegetation type is the most important variable driving the spatial pattern of SOC. According to our models, given the acidification rate in Chinese grassland soils in the future is the same as that in Chinese cropland soils during the past two decades: 0.27 and 0.48 units per 20 yr in the Inner Mongolian grasslands and the Tibetan grasslands, respectively, it will lead to 30 % and 53 % decrease in SIC in the Inner Mongolian grasslands and the Tibetan grasslands, respectively. However, negative relationship between soil pH and SOC suggests that acidification will inhibit decomposition of SOC, thus will not lead to a significant general loss of carbon from soils in these regions.

1 Introduction

The importance of soil carbon (C) in global C cycling has received considerable attention in recent decades (Schulze and Freibauer, 2005; Trumbore and Czimczik, 2008;

1870

Schlesinger and Andrews, 2000; Lal, 2004a). Soil C pool comprises two distinct components: soil organic carbon (SOC) and soil inorganic carbon (SIC). The latter includes lithogenic inorganic carbon (LIC), which comes from parent material, and pedogenic inorganic carbon (PIC), which is formed through the dissolution and precipitation of carbonate parent material. On a global scale, the estimated size of SOC in the 1-m depth is 1500 to 1600 Pg (Lal, 2004b; Monger and Gallegos, 2000; Batjes, 1996), and the size of SIC is 695 to 1738 Pg (Batjes, 1996; Eswaran et al., 1995).

C flux between the atmosphere and terrestrial ecosystems involves both SOC and SIC. The C flux between SOC pools and the atmosphere depends on biomass production, organic matter input and soil respiration (Schlesinger and Andrews, 2000; Valentini et al., 2000). SIC pools exchange C with the atmosphere through a series of physical and chemical reaction, such as C sequestration by carbonate formation or CO₂ release by acidification and leaching (Lal and Kimble, 2000; Lal, 2008; Ouyang et al., 2008). Given that global change has altered temperature, precipitation, nitrogen availability and many other environmental factors (Rockström et al., 2009; Vitousek, 1994), these changes are most likely to have great impact on soil carbon. As the largest C pool in terrestrial biosphere, even a minor change in soil carbon stocks could result in a significant alteration of atmospheric CO₂ concentration (Davidson and Janssens, 2006; Trumbore and Czimczik, 2008). Therefore, both SIC and SOC pools should be considered in order to more accurately predict future soil carbon dynamics.

In arid and semiarid regions, which cover as much as one-third of the earth's surface, SIC pools and their dynamics are important as the rate of accumulation of SIC is generally higher than in other biomes (Lal, 2008). Grasslands are one of the most widespread ecosystems in arid and semiarid regions, containing about 20 % of global soil C stocks (Jobbagy and Jackson, 2000; Scurlock and Hall, 1998; Wang and Fang, 2009). Grasslands cover more than 40 % of China's land surface, ranging from temperate grasslands in arid/semi-arid region to alpine grasslands on the Tibetan Plateau (Kang et al., 2007; Yang et al., 2010b; He et al., 2009). Because of the large spatial extent, Chinese grasslands have significant effects on regional and global carbon cycles

1871

(Ni, 2002; Yang et al., 2010b). Moreover, Chinese grasslands face rapid environmental changes (Kang et al., 2007). In recent years, many studies have been conducted to investigate the pattern, controls and dynamic of SOC (e.g., Ni, 2002; Yang et al., 2010b; Wu et al., 2003; Yu et al., 2007; Xie et al., 2007; Baumann et al., 2009) as well as SIC (e.g., Yang et al., 2010a; Mi et al., 2008; Feng et al., 2002) in Chinese grasslands. SOC stocks are reported to range from 37.7 Pg (Xie et al., 2007) to 41.0 Pg (Ni, 2002) in Chinese grasslands, and about 16.7 Pg in Northern China's grasslands (Yang et al., 2010b). On the scale of whole China, SOC densities are significantly influenced by temperature (Xie et al., 2007), while soil moisture and texture control SOC density in the Tibetan grasslands (Yang et al., 2008; Baumann et al., 2009). For SIC, total stocks of the top 1 m are estimated to vary from 53.3 Pg to 77.9 Pg (Wu et al., 2009; Mi et al., 2008; Li et al., 2007) in China. Yang et al. (2010a) estimated the SIC stock of the Tibetan grasslands to 15.2 Pg. Temperature and precipitation both show significant relationships with SIC density (Li et al., 2007; Mi et al., 2008; Yang et al., 2010a).

However, there are still some uncertainties evident in recent studies. Most of the researches focused on mean soil carbon stocks for soil at depth of 1 m or more, but little work, especially for SIC, has been devoted to investigate the patterns, controls and dynamics of soil carbon in the topsoil. The topsoil is the component of the soil system showing most rapid responses to environmental changes, such as alterations in temperature, precipitation and nitrogen deposition (Liao et al., 2009; Song et al., 2005). Changes in the topsoil are particularly important for exploring ecosystem response and functioning (Franzluebbers and Stuedemann, 2010). Moreover, recent studies spent large efforts in determining pattern, controls and dynamic of SOC, but there is less attention drawn on SIC in general (Eswaran et al., 2000; Mi et al., 2008). Consequently, we cannot simply use results of these studies to predict potential responses of soil carbon to future global changes. Further, some recent studies calculated SIC as the difference between soil total carbon (STC) and SOC (e.g., Yang et al., 2010a; Mikhailova and Post, 2006). However, the proportions of carbonates in soil total carbon are usually small (Chatterjee et al., 2009), thus this method without the direct

1872

measurement of SIC could produce a large relative error. Therefore, studies focused on SIC based on measured data help us to reduce the uncertainties of previous studies and to predict response of soil carbon to global changes.

In this study, we analyzed the topsoil data of 81 sites obtained from a regional survey across grasslands in Inner Mongolia and the Tibetan Plateau during 2006–2007. We attempt (1) to investigate the large-scale patterns of SIC and SOC in the topsoil of grasslands in Inner Mongolia and on the Tibetan Plateau, (2) to examine the changes of SIC and SOC along naturally occurring environmental gradients in order to identify key controlling factors for SIC and SOC contents, and (3) to estimate potential responses of SIC and SOC in the topsoil related to soil acidification.

2 Materials and methods

2.1 Study sites

This study was conducted in temperate grasslands in the Inner Mongolia and alpine grasslands on the Tibetan Plateau, during two expeditions in late July and early August of 2006 and 2007. We selected 81 sites along an approximately 4000 km long transect (longitude from 90.80 to 120.12° E, latitude from 30.31 to 50.19° N, and altitude from 549 to 5105 m a.s.l.) for soil and plant community sampling (Table 1, Fig. 1). Mean annual temperature (MAT) ranges from –5.8 to 4.1 °C and mean annual precipitation (MAP) ranges from 148 to 604 mm. The sites along the transect represent natural zonal grassland vegetation, including the five main vegetation types: meadow steppe, typical steppe, desert steppe, alpine steppe and alpine meadow. Field sites were selected by visual inspection of the vegetation, aiming to select sampling sites subject to minimal grazing and other anthropogenic disturbances.

1873

2.2 Soil and biomass survey

Detailed field investigations included soil profile description according to FAO (2006) and IUSS Working Group WRB (2006). Soil sampling was split into two parts: schematic soil sampling by drilling at three depth-increments (0–5, 5–10, and 10–20 cm) for SIC and SOC analyses, as well as volumetric sampling using a standard container (100 cm³ in volume) at equal depths for soil bulk density (SBD) and gravimetric water content (equalling to soil moisture, SM) determination. The sampling protocol is described in Baumann et al. (2009).

Soil samples were air-dried, all live root material was removed and the remaining soil was ground using a ball mill (NM200, Retsch, Germany). We measured soil inorganic carbon (SIC) volumetrically using an inorganic carbon analyzer (Calcimeter 08.53, Eijkelkamp, Netherland). Soil total carbon (STC) was measured by dry combustion using an elemental analyzer (VARIO EL III, Elementar, Hanau, Germany) with a combustion temperature of 1150 °C and a reduction temperature of 850 °C. Soil organic carbon (SOC) was calculated as the difference between STC and SIC. The average concentration of SIC, SOC and STC of the top 20 cm was calculated using the data from the three layers with soil bulk density, and this average value was used as site level data. Soil pH was determined in 0.01 M CaCl₂ potentiometrically. Soil total nitrogen was also determined using an elemental analyzer (PE 2400 II CHN elemental analyzer, Perkin-Elmer, Boston, Massachusetts, USA).

Additionally, we measured aboveground biomass (AGB) in three plots (1 × 1 m²) and belowground biomass (BGB) in three soil pits (0.5 × 0.5 m²) at each site. Biomass was measured by oven-drying at 60 °C to a constant weight and weighting to the nearest 0.1 g.

2.3 Climate data and statistical analyses

Climate data for each site was calculated based on linear models using latitude, longitude and altitude as predictors from 55-yr average temperature and precipitation

1874

records (1951–2005) at 680 climate stations across China (Fang et al., 2001; He et al., 2009). The calculation of potential evapotranspiration (PE) and actual evapotranspiration (AE) were based on Thornthwaite's method (Thornthwaite, 1948).

We used one-way ANOVA with Turkey's *post hoc* test to compare the effects of region, vegetation type, and soil depth on SIC, SOC and the ratio of SIC to SOC (SIC/SOC). We conducted classification and regression tree (CART) analyses to detect important variables influencing the patterns of SIC, SOC and SIC/SOC. We selected the complexity parameter as split criterion, and set the observations required for a split search at 15. Climatic factors, including MAT, MAP, growing season temperature (GST, from April to August), growing season precipitation (GSP), AE, PE, as well as biotic factors including vegetation type (VT), AGB, BGB, were used in the tree models (Table 4). Since soil pH, SM, CO₂ partial pressure and water deficiency also influenced SIC, soil pH, SM, altitude (proxy of CO₂ partial pressure) as well as the ratio of MAP to PE (MAP/PE, an index of water deficiency) were incorporated into the CART analysis when we analysed SIC and SIC/SOC. Finally, for estimating the change of SIC of the topsoil in the Inner Mongolian and Tibetan grasslands under the background of soil acidification, we built ordinary linear regression models for SIC and SIC/SOC, using the most powerful explanatory variables in the CART analysis. SIC, SOC and SIC/SOC were log transformation to achieve normal distribution (Fig. 2). All statistical analyses were performed using R 2.3.0 (R Development Core Team, 2011). The classification and regression trees were developed using the R package rpart.

3 Results

3.1 Overall patterns of SIC, SOC and SIC/SOC

Frequency distributions of SIC, SOC and SIC/SOC did not deviate significantly from log normal distributions ($P > 0.05$) neither at site level nor for each depth increment. At site level, the mean values of SIC, SOC and SIC/SOC were 0.38 %, 3.58 % and 0.15,

1875

ranging between 0.00–2.92 %, 0.32–26.34 % and 0.00–0.65, with CV of 1.36, 1.19 and 1.26, respectively (Table 3). We did not find significant differences of SIC and SIC/SOC among three soil depths ($P < 0.05$), while SOC in 0–5 cm was significantly higher than that in 10–20 cm.

At site level, topsoil SIC and SOC differed remarkably between the Inner Mongolian and Tibetan grasslands. The means of both SIC and SOC in the Tibetan grasslands (0.51 % and 5.24 %, respectively) were higher ($P < 0.01$) than those in the Inner Mongolia grasslands (0.21 % and 1.61 %). However, SIC/SOC had no significant difference between the Tibetan grasslands and the Inner Mongolian grasslands (0.18 and 0.11, $P = 0.54$). These patterns are also evident for the three soil depth increments (Table 3). Soil pH showed no significant difference between the two regions, while soils in the Tibetan grasslands generally accounted for higher soil total nitrogen and SM but lower SBD compared to the Inner Mongolian grasslands (Table 2).

On the whole, vegetation type had significant effects on SIC, SOC and SIC/SOC ($P < 0.01$). Generally, SIC concentration was higher in alpine steppe soils than in meadow steppe and typical steppe soils. SOC was higher in alpine meadow soils compared to desert steppe soils. SIC/SOC was larger as in alpine steppe soils as in the alpine meadow soils (Table 3).

3.2 Factors driving spatial variations of SIC and SOC

We developed regression tree models for SIC, SOC and SIC/SOC (Fig. 3a–c). All three trees were significantly different from a random tree ($P < 0.05$). The trees for SIC, SOC and SIC/SOC explained 69 %, 76 %, and 73 % of the variance, respectively. Tree models revealed that the spatial variations in SIC, SOC and SIC/SOC were driven by very different factors (Table 4, Fig. 3).

For SIC (Fig. 3a), the tree model had an initial split on soil pH with a threshold of 7.0, implying that soil pH was the most important variable explaining the spatial variation of SIC. As pH increases above this threshold, MAT and PE represent two thermal factors that became common to affect SIC contents, indicating a negative relationship between

1876

and Li, 1987). This would lead to higher LIC in the soil in the Tibetan grasslands. Secondly, due to higher elevation, CO₂ partial pressure is lower in the Tibetan grasslands (Körner, 2003). Lower temperature also induces lower soil respiration thus lower CO₂ partial pressure in the Tibetan grasslands soil (Kato et al., 2006). This influences the formation of pedogenic carbonate as presented in the following chemical equation:



Consequently, lower CO₂ partial pressure will move the equilibrium towards more precipitation of carbonate (Nordt et al., 2000), benefitting the formation of PIC. Although this chemical reaction is well known, no studies considered CO₂ partial pressure as an important factor in determining the large-scale pattern of SIC. For the first time, our results showed that CO₂ partial pressure may play a key role in shaping topsoil SIC in extreme high altitudinal environments.


The pattern of SOC could be attributed to climatic differences between these two regions. Low temperature leads to slower decomposition rates (Wu et al., 2003; Xie et al., 2007; Kirschbaum, 1995), and high precipitation causes high vegetation productivity (Jobbagy and Jackson, 2000; Wynn et al., 2006; Callesen et al., 2003). In addition, high moisture, and especially temporal water saturation in the Tibetan grasslands due to the frozen ground, also leads to slower decomposition rates (Baumann et al., 2009). All these factors contribute to the accumulation of SOC in the topsoil of the Tibetan grasslands.

4.2 Different controls on the large scale patterns of SIC and SOC

According to the tree models, SIC and SOC in the topsoil were affected by different factors. The pattern of SIC in the topsoil could be well explained by climate, soil physical and chemical properties (including soil pH, PE, MAT, MAP and SM), while for SOC biotic and climatic factors were predominant (including vegetation type, AGB, PE and GSP).

1879

In our research, soil pH is the most important factor for explaining the variation of SIC. Importantly, this strong relationship can be related to a decrease in HCO₃⁻ due to low pH as shown by the following equation:



Acidification induces the equilibrium towards right, thus decreasing the formation of SIC (Lal and Kimble, 2000; Suarez, 2000). In our study, 42 % of the variation in SIC could be attributed to the differences of soil pH among the 81 sites. Within different pH range (pH < 7.0 or pH > 7.0), distinct environmental factors drive the finer pattern of SIC, indicating that SIC is influenced by different processes depending on soil pH. As pH increases above the threshold, thermal factors had stronger effects than other factors, showing negative relationships with SIC in the topsoil. This is consistent with results reported in other studies (Mi et al., 2008; Yang et al., 2010a). Temperature affects CaCO₃ equilibria through its effects on CaCO₃ solubility, evapotranspiration and leaching (Lal and Kimble, 2000), but it is not appropriate to attribute variation in SIC to temperature in our research. Temperature affects biological activity especially soil respiration positively (Davidson and Janssens, 2006; Raich and Schlesinger, 1992), which increases soil CO₂ partial pressure. Moreover, In our study region, temperature decreases with increasing elevation. Consequently, higher temperature means lower altitude and thus in turn higher CO₂ partial pressure. High CO₂ partial pressure would inhibit the formation of carbonates in the topsoil (Eq. 1), leading to negative relationships between SIC and thermal factors. In the areas with pH < 7.0, SIC tends to be in a form of dissolved bicarbonate and leaching process becomes more important, consequently water condition factors like MAP and SM entered the model. However, the result of our research showing that lower MAP is corresponding to lower SIC, is in contrast to some other studies, in which SIC shows a decreasing trend along precipitation gradients (Mi et al., 2008; Nordt et al., 2000; Wang et al., 2010; Yang et al., 2010a). Further analysis suggested that most of those sites with MAP < 465 mm were distributed in the Inner Mongolia, while those sites with MAP > 465 were all located on

1880

the Tibetan Plateau. Thus the positive SIC-precipitation relationship in this study may be caused by other confounding factors that were different between the two regions.

For SOC, steppe soils present much lower SOC contents than meadow soils. As the main resource of SOC, vegetation determines quantity, quality and distribution of SOC (Jobbagy and Jackson, 2000; Poeplau et al., 2011; Quideau et al., 2001). In Chinese grasslands, meadow has higher productivity than steppe (Ma et al., 2010; Ni, 2004), which means more organic matter input into meadow soils compared to steppe sites. Moreover, alpine meadows occur under cold and humid environment. Consequently, decomposition of SOC is limited by temperature and temporal water saturation (Davidson and Janssens, 2006; Jobbagy and Jackson, 2000; Baumann et al., 2009). These two interrelated processes (i.e. high input and low decomposition) both contribute to higher SOC in the topsoil of meadow ecosystem. Within different vegetation types, finer patterns of SOC were driven by different processes. In meadows, AGB showed a strong effect on SOC in the topsoil. Because AGB reflects vegetation productivity (LeBauer and Treseder, 2008; Ni, 2004), the latter influences SOC content in the topsoil of the meadow sites by controlling the input of organic matter. When AGB increases above the threshold, PE has a negative relationship with SOC. Although high temperature may stimulate plant production as temperature is an important limiting factor for vegetation growth in the alpine meadow (Piao et al., 2006), temperature-induced higher soil decomposition overrides the resulting increase in C inputs, thus leading to a decrease of SOC. On the other hand, in steppe ecosystems, water conditions are the primary constraint to vegetation productivity (Heisler-White et al., 2008; Jobbagy et al., 2002; Sala et al., 1988). High GSP could stimulate plant production and increase soil C accumulation, thus precipitation is the most important variable in controlling SOC patterns in steppe ecosystems.

4.3 Decrease in SIC but increase in STC under soil acidification

Our study provides insights in estimating future change of SIC under the scope of soil acidification scenarios. During the past several decades, parts of Europe and Eastern

1881

North America suffered from significant acidification due to acid deposition (Bowman et al., 2008; Galloway et al., 2003). For example, in Europe soil pH decreased between 0.2 to 2 units over the past 17 to 110 yr (Stevens et al., 2009). China also has to face severe soil acidification problems due to increasing energy demand and excessive application of chemical fertilizer (Hicks et al., 2008; Larssen and Carmichael, 2000; Guo et al., 2010). Considering the important role of soil pH in controlling the pattern of SIC, soil acidification will have a great impact on topsoil's SIC stocks in the future.

Up to date, no studies have investigated the trend or extent of soil acidification in Chinese grasslands. However, soil pH records in croplands close to our study region provide circumstantial evidences to approximate the extent of soil acidification. From the 1980s to the 2000s, soil pH in croplands on the Loess Plateau and the Tibetan Plateau declined by 0.27 and 0.48 units, respectively (Zhou et al., 2009; Guo et al., 2010). Assuming the acidification rate in croplands is the same as that in grasslands and remains the same in the next 20 yr, according to our linear regression models, acidification will lead to 30 % and 53 % decrease in SIC in the Inner Mongolian and Tibetan grasslands, respectively.

Although acidification leads to decrease in SIC, carbon stock in the topsoil will not necessarily decline with soil acidification. It can be assumed, that due to accumulation of SOC, STC is likely to rise with increasing acidity, backed up by the negative relationship between soil pH and SOC as well as the even faster decline of SIC/STC compared to SIC with decreasing soil pH. This pattern can be explained by two reasons. Firstly, acidification inhibits soil microbial activities and thus the SOC decomposition rate (Francis, 1986). Secondly, N deposition, a major cause of acidification, will lead to a decrease in microbial biomass and oxidase activity (Dalmonech et al., 2010; Fisk and Fahey, 2001; Zak et al., 2008) and to an increase of SOC inputs through increasing vegetation productivity (Neff et al., 2002). Both of these will contribute to accumulation of SOC, thus leading to an increase in STC. Moreover, SIC only accounts for a very small proportion of STC on average. Therefore, soil acidification would not make a great impact on soil carbon.

1882

- Chatterjee, A., Lal, R., Wielopolski, L., Martin, M. Z., and Ebinger, M. H.: Evaluation of different soil carbon determination methods, *Crit. Rev. Plant. Sci.*, 28, 164–178, 2009.
- Chinese Academy of Sciences: Vegetation atlas of China, Science Press, Beijing, 2001.
- Dalmonech, D., Lagomarsino, A., Moscatelli, M. C., Chiti, T., and Valentini, R.: Microbial performance under increasing nitrogen availability in a Mediterranean forest soil, *Soil Biol. Biochem.*, 42, 1596–1606, 2010.
- Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature*, 440, 165–173, 2006.
- Eswaran, H., Van den Berg, E., Reich, P., and Kimble, J.: Global soil carbon resources, in: *Soils and Global Change*, edited by: Lal, R., Kimble, J. M., Levine, E., and Stewart, B. A., CRC/Lewis Publishers, Boca Raton, 27–43, 1995.
- Eswaran, H., Reich, P. F., Kimble, J. M., Beinroth, F. H., Padmanabhan, E., and Moncharoen, P.: Global carbon stocks, in: *Global Climate Change and Pedogenic Carbonates*, edited by: Lal, R., Kimble, J. M., Eswaran, H., and Stewart, B. A., CRC Press, Boca Raton, Florida, 15–25, 2000.
- Fang, J. Y., Piao, S. L., Tang, Z. Y., Peng, C. H., and Wei, J.: Interannual variability in net primary production and precipitation, *Science*, 293, 1723, 2001.
- FAO: Guidelines for soil description, 4th ed., FAO, Rome, 2006.
- Feng, Q., Endo, K. N., and Cheng, G. D.: Soil carbon in desertified land in relation to site characteristics, *Geoderma*, 106, 21–43, 2002.
- Fisk, M. C. and Fahey, T. J.: Microbial biomass and nitrogen cycling responses to fertilization and litter removal in young northern hardwood forests, *Biogeochemistry*, 53, 201–223, 2001.
- Francis, A. J.: The ecological effects of acid deposition II: acid-rain effects on soil and aquatic microbial processes, *Cell. Mol. Life Sci.*, 42, 455–465, 1986.
- Franzluebbers, A. J. and Stuedemann, J. A.: Surface soil changes during twelve years of pasture management in the Southern Piedmont USA, *Soil Sci. Soc. Am. J.*, 74, 2131–2141, 2010.
- Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., Howarth, R. W., Cowling, E. B., and Cosby, B. J.: The nitrogen cascade, *BioScience*, 53, 341–356, 2003.
- Guo, J. H., Liu, X. J., Zhang, Y., Shen, J. L., Han, W. X., Zhang, W. F., Christie, P., Goulding, K. W. T., Vitousek, P. M., and Zhang, F. S.: Significant acidification in major Chinese croplands, *Science*, 327, 1008–1010, 2010.
- He, J.-S., Wang, X., Flynn, D. F. B., Wang, L., Schmid, B., and Fang, J. Y.: Taxonomic, phyloge-

1885

- netic, and environmental trade-offs between leaf productivity and persistence, *Ecology*, 90, 2779–2791, 2009.
- Heisler-White, J. L., Knapp, A. K., and Kelly, E. F.: Increasing precipitation event size increases aboveground net primary productivity in a semi-arid grassland, *Oecologia*, 158, 129–140, 2008.
- Hicks, W. K., Kuylentierna, J. C. I., Owen, A., Dentener, F., Seip, H. M., and Rodhe, H.: Soil sensitivity to acidification in Asia: status and prospects, *Ambio*, 37, 295–303, 2008.
- IUSS Working Group WRB: World reference base for soil resources 2006, *World Soil Resources Reports*, No. 103, FAO, Rome, 2006.
- Jobbagy, E. G. and Jackson, R. B.: The vertical distribution of soil organic carbon and its relation to climate and vegetation, *Ecol. Appl.*, 10, 423–436, 2000.
- Jobbagy, E. G., Sala, O. E., and Paruelo, J. M.: Patterns and controls of primary production in the Patagonian steppe: a remote sensing approach, *Ecology*, 83, 307–319, 2002.
- Körner, C.: *Alpine plant life: functional plant ecology of high mountain ecosystems*, Springer Berlin, Heidelberg, 2003.
- Kang, L., Han, X. G., Zhang, Z. B., and Sun, O. J.: Grassland ecosystems in China: review of current knowledge and research advancement, *Philos. T. R. Soc. B*, 362, 997–1008, 2007.
- Kato, T., Tang, Y., Gu, S., Hirota, M., Du, M., Li, Y., and Zhao, X.: Temperature and biomass influences on interannual changes in CO₂ exchange in an alpine meadow on the Qinghai-Tibetan Plateau, *Glob. Change Biol.*, 12, 1285–1298, 2006.
- Kirschbaum, M. U. F.: The temperature dependence of soil organic matter decomposition, and the effect of global warming on soil organic C storage, *Soil Biol. Biochem.*, 27, 753–760, 1995.
- Lal, R.: Soil carbon sequestration impacts on global climate change and food security, *Science*, 304, 1623–1627, 2004a.
- Lal, R.: Agricultural activities and the global carbon cycle, *Nutr. Cycl. Agroecosys.*, 70, 103–116, 2004b.
- Lal, R.: Carbon sequestration, *Philos. T. R. Soc. B*, 363, 815–830, 2008.
- Lal, R. and Kimble, J. M.: Pedogenic carbonates and the global carbon cycle, in: *Global Climate Change and Pedogenic Carbonates*, edited by: Lal, R., Kimble, J. M., Eswaran, H., and Stewart, B. A., CRC Press, Boca Raton, Florida, 1–14, 2000.
- Larssen, T. and Carmichael, G. R.: Acid rain and acidification in China: the importance of base cation deposition, *Environ. Pollut.*, 110, 89–102, 2000.

1886

- LeBauer, D. S. and Treseder, K. K.: Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed, *Ecology*, 89, 371–379, 2008.
- Li, Z. P., Han, F. X., Su, Y., Zhang, T. L., Sun, B., Monts, D. L., and Plodinec, M. J.: Assessment of soil organic and carbonate carbon storage in China, *Geoderma*, 138, 119–126, 2007.
- 5 Liao, Q. L., Zhang, X. H., Li, Z. P., Pan, G. X., Smith, P., Jin, Y., and Wu, X. M.: Increase in soil organic carbon stock over the last two decades in China's Jiangsu Province, *Glob. Change Biol.*, 15, 861–875, 2009.
- Ma, W. H., He, J.-S., Yang, Y. H., Wang, X. P., Liang, C. Z., Anwar, M., Zeng, H., Fang, J. Y., and Schmid, B.: Environmental factors covary with plant diversity-productivity relationships among Chinese grassland sites, *Glob. Ecol. Biogeogr.*, 19, 233–243, 2010.
- 10 Mi, N., Wang, S. Q., Liu, J. Y., Yu, G. R., Zhang, W. J., and Jobbaagy, E.: Soil inorganic carbon storage pattern in China, *Glob. Change Biol.*, 14, 2380–2387, 2008.
- Mikhailova, E. A. and Post, C. J.: Effects of land use on soil inorganic carbon stocks in the Russian Chernozem, *J. Environ. Qual.*, 35, 1384–1388, 2006.
- 15 Monger, H. C. and Gallegos, R. A.: Biotic and abiotic processes and rates of pedogenic carbonate accumulation, in: *Global Climate Change and Pedogenic Carbonates*, edited by: Lal, R., Kimble, J. M., Eswaran, H., and Stewart, B. A., CRC Press, Boca Raton, Florida, 2000.
- Neff, J. C., Townsend, A. R., Gleixner, G., Lehman, S. J., Turnbull, J., and Bowman, W. D.: Variable effects of nitrogen additions on the stability and turnover of soil carbon, *Nature*, 419, 915–917, 2002.
- 20 Ni, J.: Carbon storage in grasslands of China, *J. Arid. Environ*, 50, 205–218, 2002.
- Ni, J.: Estimating net primary productivity of grasslands from field biomass measurements in temperate northern China, *Plant Ecol.*, 174, 217–234, 2004.
- Nordt, L. C., Wilding, L. P., and Drees, L. R.: Pedogenic carbonate transformations in leaching soil system: implication for the global C cycle, in: *Global Climate Change and Pedogenic Carbonates*, edited by: Lal, R., Kimble, J. M., Eswaran, H., and Stewart, B. A., CRC Press, Boca Raton, Florida, 43–64, 2000.
- 25 Ouyang, X. J., Zhou, G. Y., Huang, Z. L., Liu, J. X., Zhang, D. Q., and Li, J.: Effect of simulated acid rain on potential carbon and nitrogen mineralization in forest soils, *Pedosphere*, 18, 503–514, 2008.
- 30 Piao, S. L., Fang, J. Y., and He, J. -S.: Variations in vegetation net primary production in the Qinghai-Xizang Plateau, China, from 1982 to 1999, *Clim. Change*, 74, 253–267, 2006.
- Poepflau, C., Don, A., Vesterdal, L., Leifeld, J., Van Wesemael, B. A. S., Schumacher, J., and

1887

- Gensior, A.: Temporal dynamics of soil organic carbon after land-use change in the temperate zone: carbon response functions as a model approach, *Glob. Change Biol.*, 17, 2415–2427, 2011.
- 5 Quideau, S. A., Chadwick, O. A., Benesi, A., Graham, R. C., and Anderson, M. A.: A direct link between forest vegetation type and soil organic matter composition, *Geoderma*, 104, 41–60, 2001.
- R Development Core Team: R: a language and environment for statistical computing, R Foundation for Statistical Computing Vienna, Austria, 2011.
- 10 Raich, J. W. and Schlesinger, W. H.: The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate, *Tellus B*, 44, 81–99, 1992.
- Rockström, J., Steffen, W., Noone, K., Persson, A., Chapin, F. S., Lambin, E. F., Lenton, T. M., Scheffer, M., Folke, C., Schellnhuber, H. J., Nykvist, B., de Wit, C. A., Hughes, T., van der Leeuw, S., Rodhe, H., Sorlin, S., Snyder, P. K., Costanza, R., Svedin, U., Falkenmark, M., Karlberg, L., Corell, R. W., Fabry, V. J., Hansen, J., Walker, B., Liverman, D., Richardson, K., Crutzen, P., and Foley, J. A.: A safe operating space for humanity, *Nature*, 461, 472–475, 2009.
- 15 Sala, O. E., Parton, W. J., Joyce, L. A., and Lauenroth, W. K.: Primary production of the central grassland region of the United States, *Ecology*, 69, 40–45, 1988.
- Schlesinger, W. H. and Andrews, J. A.: Soil respiration and the global carbon cycle, *Biogeochemistry*, 48, 7–20, 2000.
- 20 Schulze, E. D. and Freibauer, A.: Carbon unlocked from soils, *Nature*, 437, 205–206, 2005.
- Scurlock, J. M. O. and Hall, D. O.: The global carbon sink: a grassland perspective, *Glob. Change Biol.*, 4, 229–233, 1998.
- Song, G. H., Li, L. Q., Pan, G. X., and Zhang, Q.: Topsoil organic carbon storage of China and its loss by cultivation, *Biogeochemistry*, 74, 47–62, 2005.
- 25 Stevens, C. J., Dise, N. B., and Gowing, D. J.: Regional trends in soil acidification and exchangeable metal concentrations in relation to acid deposition rates, *Environ. Pollut.*, 157, 313–319, 2009.
- Suarez, D. L.: Impact of agriculture on CO₂ as affected by changes in inorganic carbon, in: *Global Climate Change and Pedogenic Carbonates*, edited by: Lal, R., Kimble, J. M., Eswaran, H., and Stewart, B. A., CRC Press, Boca Raton, Florida, 257–272, 2000.
- 30 Thornthwaite, C. W.: An approach toward a rational classification of climate, *Geograph. Rev.*, 38, 55–94, 1948.

1888

- Trumbore, S. E. and Czimczik, C. I.: An uncertain future for soil carbon, *Science*, 321, 1455–1456, 2008.
- Valentini, R., Matteucci, G., Dolman, A. J., Schulze, E. D., Rebmann, C., Moors, E. J., Granier, A., Gross, P., Jensen, N. O., Pilegaard, K., Lindroth, A., Grelle, A., Bernhofer, C., Grunwald, T., Aubinet, M., Ceulemans, R., Kowalski, A. S., Vesala, T., Rannik, U., Berbigier, P., Loustau, D., Guomundsson, J., Thorgeirsson, H., Ibrom, A., Morgenstern, K., Clement, R., Moncrieff, J., Montagnani, L., Minerbi, S., and Jarvis, P. G.: Respiration as the main determinant of carbon balance in European forests, *Nature*, 404, 861–865, 2000.
- Vitousek, P. M.: Beyond global warming: ecology and global change, *Ecology*, 75, 1861–1876, 1994.
- Wang, W. and Fang, J.: Soil respiration and human effects on global grasslands, *Global Planet. Change*, 67, 20–28, 2009.
- Wang, Y. G., Li, Y., Ye, X. H., Chu, Y., and Wang, X. P.: Profile storage of organic/inorganic carbon in soil: from forest to desert, *Sci. Total Environ.*, 408, 1925–1931, 2010.
- Wu, H. B., Guo, Z. T., and Peng, C. H.: Distribution and storage of soil organic carbon in China, *Global Biogeochem. Cy.*, 17, 1048, doi:10.1029/2001GB001844, 2003.
- Wu, H. B., Guo, Z. T., Gao, Q., and Peng, C. H.: Distribution of soil inorganic carbon storage and its changes due to agricultural land use activity in China, *Agr. Ecosys. Environ.*, 129, 413–421, 2009.
- Wynn, J. G., Bird, M. I., Vellen, L., Grand-Clement, E., Carter, J., and Berry, S. L.: Continental-scale measurement of the soil organic carbon pool with climatic, edaphic, and biotic controls, *Global Biogeochem. Cy.*, 20, GB1007, doi:10.1029/2005GB002576, 2006.
- Xie, Z. B., Zhu, J. G., Liu, G., Cadisch, G., Hasegawa, T., Chen, C. M., Sun, H. F., Tang, H. Y., and Zeng, Q.: Soil organic carbon stocks in China and changes from 1980s to 2000s, *Glob. Change Biol.*, 13, 1989–2007, 2007.
- Xiong, Y., and Li, Q. D.: *Soils of China*, Science Press, Beijing, 1987 (in Chinese).
- Yang, Y. H., Fang, J. Y., Tang, Y. H., Ji, C. J., Zheng, C. Y., He, J. S., and Zhu, B. A.: Storage, patterns and controls of soil organic carbon in the Tibetan grasslands, *Glob. Change Biol.*, 14, 1592–1599, doi:10.1111/j.1365-2486.2008.01591.x, 2008.
- Yang, Y. H., Fang, J. Y., Ji, C. J., Ma, W. H., Su, S. S., and Tang, Z. Y.: Soil inorganic carbon stock in the Tibetan alpine grasslands, *Global Biogeochem. Cy.*, 24, GB4022, doi:10.1029/2010GB003804, 2010a.
- Yang, Y. H., Fang, J. Y., Ma, W. H., Smith, P., Mohammad, A., Wang, S. P., and Wang, W.: Soil

1889

- carbon stock and its changes in northern China's grasslands from 1980s to 2000s, *Glob. Change Biol.*, 16, 3036–3047, 2010b.
- Yu, D. S., Shi, X. Z., Wang, H., Sun, W. X., Chen, J. M., Liu, Q. H., and Zhao, Y. C.: Regional patterns of soil organic carbon stocks in China, *J. Environ. Manage.*, 85, 680–689, 2007.
- Zak, D. R., Holmes, W. E., Burton, A. J., Pregitzer, K. S., and Talhelm, A. F.: Simulated atmospheric NO_3^- deposition increases soil organic matter by slowing decomposition, *Ecol. Appl.*, 18, 2016–2027, 2008.
- Zhou, Z., Du, S., and Liu, G.: Acidification of surface soil in croplands in the semiarid middle Tibet Plateau, China, 2009 International Conference on Environmental Science and Information Application Technology, 2009, 209–212, 2009.

1890

Table 1. Description of the study regions. Mean annual temperature (MAT), mean annual precipitation (MAP), growing season temperature (GST), growing season precipitation (GSP), averaged topsoil pH, soil bulk density (SBD), soil moisture (SM) and soil total nitrogen (STN) of the sampling sites are shown.

	Overall	Inner Mongolian grasslands	Tibetan grasslands
No. of site	81	36	45
Longitude (° E)	90.80–120.12	111.83–120.12	90.80–101.48
Latitude (° N)	30.31–50.19	41.79–50.19	30.31–37.28
Altitude (m)	549–5105	549–1418	2925–5105
MAT (°C)	–5.8–4.1	–2.6–4.1	–5.8–2.6
GST (°C)	1.5–16.9	11.2–16.9	1.5–11
MAP (mm yr ⁻¹)	148–604	148–436	218–604
GSP (mm yr ⁻¹)	115–402	115–343	133–402
Soil pH	5.2–8.2	5.7–8.2	5.2–7.6
SBD (g cm ⁻³)	0.25–1.63	0.94–1.63	0.25–1.43
SM (V/V %)	0.73–68.11	2.04–16.10	0.73–68.11
STN (g/g %)	0.04–1.52	0.05–0.45	0.04–1.52

1891

Table 2. Topsoil properties in the Inner Mongolian and Tibetan grasslands across sampling sites. Different letters indicate statistical significance at $P < 0.05$. SBD = soil bulk density; SM = soil moisture; STN = soil total nitrogen; CV = coefficient of variation.

		Inner Mongolian grasslands	Tibetan grasslands
Soil pH	<i>n</i>	36	45
	Mean	6.9 a	6.8 a
	CV	0.10	0.09
	Range	5.7–8.2	5.2–7.6
SBD (g cm ⁻³)	<i>n</i>	36	45
	Mean	1.3 b	0.95 a
	CV	0.13	0.33
	Range	0.94–1.63	0.25–1.43
SM (V/V %)	<i>n</i>	32	44
	Mean	7.29 a	20.95 b
	CV	0.55	0.85
	Range	2.04–16.10	0.73–68.11
STN (g/g %)	<i>n</i>	36	45
	Mean	0.17 a	0.44 b
	CV	0.54	0.81
	Range	0.05–0.45	0.04–1.52

1892

Table 3. Statistical description of soil inorganic carbon (SIC), soil organic carbon (SOC) content and SIC to soil total carbon ratio (SIC/STC) in each depth increment in different regions and vegetation types. Different letters indicate statistical significance at $P < 0.05$. CV = coefficient of variation.

		SIC (%)				SOC (%)				SIC/STC			
		0–5 cm	5–10 cm	10–20 cm	Site level	0–5 cm	5–10 cm	10–20 cm	Site level	0–5 cm	5–10 cm	10–20 cm	Site level
Overall	Mean	0.31	0.35	0.44	0.38	4.78	3.74	2.92	3.58	0.11	0.13	0.18	0.15
	CV	1.49	1.52	1.33	1.36	1.08	1.17	1.32	1.19	1.46	1.43	1.16	1.26
	Range	0.00–1.92	0.00–2.28	0.00–2.61	0.00–2.20	0.29–27.51	0.30–25.43	0.28–26.00	0.32–26.34	0.00–0.66	0.00–0.66	0.00–0.74	0.00–0.65
	<i>n</i>	87	87	87	81	87	87	87	81	87	87	87	81
Region	Mean	0.14a	0.18a	0.25a	0.22a	2.23a	1.58a	1.26a	1.61a	0.07a	0.09a	0.14a	0.11a
Inner Mongolian grasslands	CV	2.09	2.30	1.81	1.89	0.72	0.66	0.65	0.66	1.44	1.66	1.31	1.39
	Range	0.01–1.59	0.00–2.12	0.01–2.37	0.01–2.10	0.29–7.54	0.30–4.42	0.28–3.60	0.32–4.55	0.00–0.39	0.00–0.53	0.00–0.65	0.00–0.54
	<i>n</i>	39	39	39	36	39	39	36	36	39	39	39	36
Tibetan grasslands	Mean	0.45b	0.49b	0.58b	0.51b	6.84b	5.50b	4.27b	5.15b	0.15a	0.17a	0.21a	0.18a
	CV	1.18	1.19	1.07	1.10	0.89	0.95	1.11	0.99	1.30	1.27	1.06	1.16
	Range	0.00–1.92	0.00–2.28	0.00–2.61	0.00–2.20	0.41–27.51	0.35–25.43	0.45–26.00	0.45–26.34	0.00–0.66	0.00–0.66	0.00–0.74	0.00–0.65
	<i>n</i>	48	48	48	45	48	48	48	45	48	48	48	45
Vegetation type	Mean	0.06ab	0.04ab	0.16ab	0.11a	3.21b	2.46b	2.04b	2.43b	0.03a	0.02a	0.09ab	0.06ab
Meadow steppe	CV	0.88	0.47	1.71	1.41	0.54	0.57	0.53	0.53	1.21	0.68	1.49	1.46
	Range	0.02–0.19	0.02–0.07	0.01–0.78	0.01–0.44	0.88–5.29	0.73–4.17	0.59–3.60	0.70–4.15	0.00–0.11	0.00–0.05	0.00–0.37	0.00–0.25
	<i>n</i>	7	7	7	7	7	7	7	7	7	7	7	
Typical steppe	Mean	0.16a	0.21a	0.28a	0.24a	2.42b	1.64ab	1.25ab	1.65ab	0.05a	0.08a	0.12ab	0.09ab
	CV	2.23	2.36	1.93	2.02	0.64	0.52	0.51	0.55	1.81	1.79	1.46	1.52
	Range	0.01–1.59	0.00–2.12	0.01–2.37	0.01–2.10	0.69–7.54	0.57–4.418	0.48–3.51	0.59–4.55	0.00–0.39	0.00–0.50	0.01–0.63	0.01–0.51
	<i>n</i>	24	24	24	22	24	24	22	22	24	24	24	22
Desert steppe	Mean	0.23ab	0.31abc	0.44ab	0.39ab	1.07a	0.98a	0.80a	0.98a	0.17b	0.20ab	0.26ab	0.24bc
	CV	1.08	1.19	1.11	1.00	0.89	0.82	0.75	0.73	0.81	0.95	0.75	0.79
	Range	0.01–0.76	0.01–0.92	0.02–1.25	0.01–0.94	0.29–3.40	0.30–2.96	0.28–2.07	0.32–2.62	0.02–0.45	0.02–0.53	0.02–0.65	0.02–0.54
	<i>n</i>	10	10	10	9	10	10	9	10	10	10	9	
Alpine steppe	Mean	0.71c	0.76c	0.83b	0.78b	2.42b	2.07ab	1.80b	1.97ab	0.29b	0.30b	0.34b	0.32c
	CV	0.82	0.90	0.89	0.88	0.72	0.63	0.61	0.65	0.75	0.77	0.68	0.69
	Range	0.00–1.92	0.04–2.28	0.01–2.61	0.02–2.20	0.41–6.24	0.35–4.33	0.45–4.00	0.45–4.19	0.00–0.66	0.01–0.66	0.01–0.74	0.01–0.65
	<i>n</i>	18	18	18	17	18	18	18	18	18	18	18	17
Alpine meadow	Mean	0.27b	0.31bc	0.39ab	0.30ab	10.02c	7.94c	6.03c	7.47c	0.05a	0.08a	0.11a	0.07a
	CV	1.63	1.51	1.27	1.25	0.61	0.71	0.92	0.76	2.01	2.01	1.53	1.66
	Range	0.00–1.698	0.00–1.75	0.00–1.71	0.00–1.68	1.07–27.51	0.82–25.43	0.69–26.00	0.81–26.34	0.00–0.40	0.00–0.63	0.00–0.67	0.00–0.55
	<i>n</i>	28	28	28	26	28	28	28	26	28	28	28	26

1893

Table 4. Variables included and selected in the regression tree analysis. Alt = altitude; MAT = mean annual temperature; GST = growing season temperature; MAP = mean annual precipitation; GSP = growing season precipitation; PE = potential evapotranspiration; AE = actual evapotranspiration; AP/PE = the ratio of MAP to PE; VT = vegetation type; AGB = above-ground biomass; BGB = belowground biomass; SBD = soil bulk density; SM = soil moisture.

Variable	<i>n</i>	Mean	SD	Variables used to build the tree models			Variables selected finally		
				SIC	SOC	SIC/STC	SIC	SOC	SIC/STC
Alt	80	2692	1692.4	Yes		Yes			
MAT	79	-1.1	2.42	Yes	Yes	Yes	Yes	Yes	
GST	75	8.8	4.63	Yes	Yes	Yes			
MAP	79	384.0	113.66	Yes	Yes	Yes	Yes		
GSP	75	270.5	63.01	Yes	Yes	Yes		Yes	
PE	75	432.0	92.30	Yes	Yes	Yes	Yes	Yes	
AE	75	321.7	59.81	Yes	Yes	Yes			
AP/PE	74	1.0	0.43	Yes		Yes			
VT	81	-	-	Yes	Yes	Yes		Yes	
AGB	77	96.4	59.41	Yes	Yes	Yes		Yes	
BGB	73	113.3	1515.15	Yes	Yes	Yes			
pH	81	6.88	0.563	Yes		Yes	Yes	Yes	
SBD	81	1.11	0.315	Yes					
SM	76	15.2	15.27	Yes		Yes	Yes	Yes	

1894




Fig. 1. Vegetation map of Chinese grasslands and location of sampling sites (1 : 1 000 000) (Chinese Academy of Sciences, 2001).

1895




Fig. 2. Frequency distributions of SIC (a–d), SOC (e–h) and SIC/STC (i–l) in Chinese grasslands across all sampling sites at soil depths of 0–5 cm (a, e, i), 5–10 cm (b, f, j), 10–20 cm (c, g, k) and at site level (d, h, l). All distributions have no significant differences compared with normal distribution at $P < 0.05$.

1896

